

The Huygens principle and cosmological gravitational waves in the Regge-Wheeler gauge.

Edward Malec and Grzegorz Wylęzek

Institute of Physics, Jagiellonian University, 30-059 Cracow, Reymonta 4, Poland.

We study the propagation of axial gravitational waves in Friedman universes. The evolution equation is obtained in the Regge-Wheeler gauge. The gravitational waves obey the Huygens principle in the radiation dominated era, but in the matter dominated universe their propagation depends on their wavelengths, with the scale fixed essentially by the Hubble radius. Short waves practically satisfy the Huygens principle while long waves can backscatter off the curvature of a spacetime.

I. INTRODUCTION.

Information about the topological type of the Universe can be obtained through the analysis of the Hubble relation [1]. One of the key and well known facts underlying derivation of the Hubble law is that electromagnetic waves propagate along null cones, that is that they satisfy the Huygens principle in Friedman universes [2]. It is known that gravitational waves do not obey the Huygens principle – they can backscatter on the curvature of the spacetime and a part of the radiation would come with a delay. The observation of the effects of the backscatter opens, in principle at least, a new local way to see the global topology in a way analogous to effects known in the Schwarzschild spacetime ([3] – [5]).

The main goal of this paper is to explain whether and when these effects can manifest. We consider gravitational waves propagating in a fixed Friedman universe. The gravitational waves correspond to the so-called axial modes and they fulfill the Regge-Wheeler gauge condition [6]. The propagation equations become particularly simple in this gauge – they reduce to a single equation. This analysis with the Regge-Wheeler condition is novel in the cosmological literature, up to our knowledge. The form of the equation immediately implies that axial modes do satisfy the Huygens principle in the radiation epoch. On the other hand, the principle might well be broken in the matter-dominated epoch, depending on the wavelength of the gravitational radiation. The wavelength scale λ is given by the average energy density ρ , $\lambda \propto 1/\sqrt{\rho}$; under suitable conditions λ is of the order of the Hubble radius – that is, of the radius of the visible part of the universe. The effects of backscattering are negligible in the short-range spectrum with wavelengths much smaller than λ and they can possibly manifest for longer waves. That would mean that eventual effects cannot be observed by detectors of gravitational waves but they might be responsible for some superhorizon perturbations.

II. MAIN EQUATIONS.

Let $ds^2 = a^2 (-d\eta^2 + dr^2 + \sigma^2 d\Omega^2)$ be the line element of a Friedman cosmological model. Here η is the conformal time and the coordinate radius $\sigma = r, \sin r$ and $\sinh r$ for the flat ($k = 0$), closed ($k = 1$) and open ($k = -1$) universes, respectively. The axial perturbations of this metric in the Regge-Wheeler gauge [6] are defined as follows

$$\begin{aligned} g_{0\phi} &= h_0(\eta, r) \sin \theta \partial_\theta Y_{l0} \\ g_{1\phi} &= h_1(\eta, r) \sin \theta \partial_\theta Y_{l0}; \end{aligned} \quad (1)$$

strictly saying, this a ϕ -independent subset of a larger class, but the general axial modes conform to the same dynamical equations as the ones displayed below. The linearized Einstein equations (for $l \geq 2$, applying arguments of [6] and [7]) are

$$\partial_\eta h_0 = \partial_r h_1, \quad (2)$$

$$\partial_\eta^2 h_1 - \partial_\eta \partial_r h_0 + 2\partial_\eta h_0 \frac{\partial_r \sigma}{\sigma} - 2\partial_\eta \left(h_1 \frac{\partial_\eta a}{a} \right) + [l(l+1) - 2] \frac{h_1}{\sigma^2} = 0, \quad (3)$$

$$\partial_r^2 h_0 - \partial_\eta \partial_r h_1 - 2\partial_\eta h_1 \frac{\partial_r \sigma}{\sigma} + 2 \frac{\partial_\eta a}{a\sigma^2} \partial_r (\sigma^2 h_1) - l(l+1) \frac{h_0}{\sigma^2} + 4kh_0 = 0. \quad (4)$$

This is a system of three equations on the two unknown functions h_0 and h_1 , but it is not over-determined, as we demonstrate in what follows. Acting on both sides of (4) by the operator $\sigma^2 \partial_\eta$ and using (2) in order to eliminate $\partial_\eta h_0$ in the obtained expression, one arrives eventually at

$$\partial_r F = 4k\sigma^2 \partial_r h_1 - 2(\partial_r^2 h_1) \sigma \partial_r \sigma. \quad (5)$$

Here F is given

$$F = \sigma^2 (\partial_\eta^2 h_1 - \partial_r^2 h_1) + l(l+1) h_1 - 2\sigma^2 \partial_\eta \left(h_1 \frac{\partial_\eta a}{a} \right). \quad (6)$$

Analogously, replacing $\partial_\eta h_0$ by $\partial_r h_1$ – thanks to (2) – in Eq. (3), one can obtain

$$F = -2\sigma \partial_r \sigma \partial_r h_1 + 2h_1. \quad (7)$$

Comparison of (5) and (7) allows one to conclude, that the two equations are equivalent – modulo terms in h_1 which are r -independent – if the compatibility condition

$$-2\partial_r h_1 [\partial_r (\sigma \partial_r \sigma) - 1 + 2k\sigma^2] = 0. \quad (8)$$

is satisfied. It is a straightforward exercise to check out that the expression within the square bracket vanishes for all Friedman models. If one assumes in addition compact support of initial data, then the above mentioned ambiguity disappears and one can conclude that the two equations, (5) and (7), are equivalent. The whole dynamical content of the evolution is described by Eqs. (6) and (7),

$$\sigma^2 (\partial_\eta^2 h_1 - \partial_r^2 h_1) + l(l+1) h_1 - 2\sigma^2 \partial_\eta \left(h_1 \frac{\partial_\eta a}{a} \right) = -2\sigma \partial_r \sigma \partial_r h_1 + 2h_1. \quad (9)$$

Insert now $Q = h_1/(\sigma a)$ into equation (9); this yields

$$\partial_\eta^2 Q - \partial_r^2 Q + \frac{l(l+1)}{\sigma^2} Q - kQ - \frac{\partial_\eta^2 a}{a} Q = 0. \quad (10)$$

One can show, using Friedman equations, that in the radiation epoch $\partial_\eta^2 a = -ka$ while in the dust epoch $\partial_\eta^2 a = -ka + 4\pi\rho a^3/3$. Thence the gravitational wave evolves during radiation era according to the equation

$$\partial_\eta^2 Q - \partial_r^2 Q + \frac{l(l+1)}{\sigma^2} Q = 0. \quad (11)$$

During the matter (and, in particular, dust) dominated epoch, in turn, the gravitational waves satisfy

$$\partial_\eta^2 Q - \partial_r^2 Q + \frac{l(l+1)}{\sigma^2} Q - \frac{4\pi\rho a^2}{3} Q = 0. \quad (12)$$

III. ANALYSIS.

During the radiation epoch the Huygens principle holds true. Indeed, the form of the equation (11) is exactly that of the electromagnetic fields in Friedman spacetimes. Therefore its general solution has the form (see, for instance, [8])

$$\phi_l(r, \eta) = \sigma^l \underbrace{\partial_r \frac{1}{\sigma} \partial_r \frac{1}{\sigma} \cdots \partial_r \left(\frac{f+g}{\sigma} \right)}_l \quad (13)$$

where the free functions f and g depend on the combinations $r - \eta$ or $r + \eta$, respectively. Clearly, from this form of the solution, compact pulses of radiation have to move within space-time regions bounded by a pair of null cones $\eta = \pm r + \text{const}$. In particular shocks have to move along null cones.

Now we shall discuss the evolution of gravitational waves in the matter-dominated era. Assume initial data of compact support $\sigma(r) \leq \sigma(r_0)$ at the beginning η_0 of this epoch. Thus at a time η the radius of the support of Q is

bounded from above by $\sigma(r_0 + \eta)$; assume in addition $\sigma(r + \eta) < \pi/2$ and $k/a^2 \ll H^2$ in the case of the $k = 1$ universe. Let $R(\eta) = a(\eta)\sigma(r + \eta)$ be the related areal radius. Define the critical-length parameter $\lambda(\eta) \equiv \sqrt{3l(l+1)/(4\pi\rho)}$. Let us remark that, in the case of the flat universe, $\lambda(\eta)$ is of the order of the quantity $\sqrt{2l(l+1)}/H$ (where H is the Hubble constant). This is related to the Hubble radius and for low value of l is of the order of the radius of the visible part of the universe. In the case of closed and open universes $\lambda(\eta)$ can be bounded from above or below, respectively, by $\sqrt{2l(l+1)}/H$ – again, for small l , being close to the Hubble radius $1/H$. One can write, taking into account that the product $\rho(\eta)a^3(\eta)$ is constant during the matter dominated era, a sequence of inequalities relating the ratio $N(\eta)$ of the two radii, taken at successive moments of conformal time,

$$N(\eta_1) \equiv \frac{R^2(\eta_1)}{\lambda^2(\eta_1)} \leq \frac{R^2(\eta_2)}{\lambda^2(\eta_2)} \frac{a(\eta_2)}{a(\eta_1)} = N(\eta_2) (1 + z(1, 2)). \quad (14)$$

Here $z(1, 2) \equiv \frac{a(\eta_2)}{a(\eta_1)}$ is the cosmological redshift and $\eta_0 \leq \eta_1 < \eta_2$.

The ratio of the last two terms of Eq. (12) is bounded from above by $4\pi\rho a^2(\eta)\sigma^2(r + \eta)/(3l(l+1))$. This can be expressed as $N(\eta) = R^2(\eta)/\lambda^2(\eta)$. Clearly, if $N(\eta_2)(1 + z(1, 2)) < 1$, then $N(\eta_1) < 1$. That suggests that for $\eta \in (\eta_0, \eta_2)$ the evolution is well described by the equation (11) and the solution is close to that given by (13). Therefore there is no backscatter. And conversely, if the above product is at least of the order of unity, then there can exist a backscattered part, with characteristics possibly allowing the identification of the type of the universe. This heuristic reasoning can be made rigorous, as outlined below.

One can easily check that the energy

$$E(Q) \equiv \frac{1}{2} \int_0^\infty dr \left[(\partial_\eta Q)^2 + (\partial_r Q)^2 + \frac{l(l+1)}{\sigma^2} Q^2 \right] \quad (15)$$

is conserved by the evolution equation (12). Split the sought solution of (12) as $Q = Q_0 + x$, where Q_0 satisfies Eq. (11) and initially $Q = Q_0, \partial_\eta Q = \partial_\eta Q_0$. Thence x solves the equation

$$\partial_\eta^2 x - \partial_r^2 x + \frac{l(l+1)}{\sigma^2} x = \frac{4\pi\rho a^2}{3} (x + Q_0) \quad (16)$$

and satisfies homogeneous initial conditions $x = \partial_\eta x = 0$. The energy $E(x)$ of the field x satisfies the equation

$$\frac{d}{d\eta} E(x) = \int_0^\infty dr \frac{4\pi\rho a^2}{3} (Q_0 + x) \partial_\eta x. \quad (17)$$

One notices that $\sigma(r_0 + \eta) \geq \sigma(r)$ for all points r lying within the support of Q_0 .

Therefore equation (17) implies the inequality

$$\frac{d}{d\eta} E(x) \leq \frac{4\pi\rho a^2}{3\sqrt{l(l+1)}} \sigma(r_0 + \eta) \int_0^\infty dr \frac{\sqrt{l(l+1)}}{\sigma(r)} |(Q_0 + x) \partial_\eta x|. \quad (18)$$

Now, making use of the Schwarz inequality and bounding the relevant factors on the the right hand side of (18) by energies $E(x)$ and $E(Q_0)$, dividing both sides by $\sqrt{E(x)}$ and rearranging the resulting inequality, one obtains

$$e^{\alpha(\eta)} \frac{d}{d\eta} \left(e^{-\alpha(\eta)} \sqrt{E(x, \eta)} \right) \leq \frac{d\alpha}{d\eta} \sqrt{E(Q_0)}, \quad (19)$$

where

$$\alpha = \int_{\eta_0}^\eta d\tilde{\eta} \frac{4\pi\rho a^2}{3\sqrt{l(l+1)}} \sigma(r_0 + \tilde{\eta}). \quad (20)$$

In the case of flat and open universes one has $4\pi\rho a^2/3 \leq H^2 a^2/2$, while for the closed universe there is an approximate equality $4\pi\rho a^2/3 \approx H^2 a^2/2$, under the condition imposed earlier. Notice now that $d\eta H a^2 = da$; change the integration variable and observe that $\sigma(r_0 + \eta)$ and $H(\eta)$ are increasing with η , in order to get the estimate

$$\alpha \leq \sigma(r_0 + \eta)(a(\eta) - a(\eta_0)) \frac{H(\eta)}{2\sqrt{l(l+1)}}. \quad (21)$$

This can be written concisely in terms of the areal radius and the critical parameter,

$$\alpha \leq \beta \equiv \frac{R(\eta)}{\lambda(\eta)\sqrt{2}}. \quad (22)$$

Finally, one arrives at the inequality

$$E(x, \eta) \leq E(Q_0)(e^\beta - 1)^2. \quad (23)$$

Similar estimates can be obtained for the energy $E(\partial_r x)$ and of higher derivatives of x . Employing these, one can use standard Sobolev estimates [9] in order to show that the field x (and its derivatives, if needed) can be pointwise small, under the condition that initial data at the time $\eta_0 = 0$ have compact support and that $R(\eta)/(\lambda(\eta)) \ll 1$.

Let us remark that we omit boundary terms coming from $r = 0$, without discussing subtleties related with the behaviour of solutions at $r = 0$, when integrating by parts. This is justified by the fact that "good" initial data of our $1+1$ evolution problem – i. e., those inherited from the initial $1+3$ formulation of the evolution of gravitational waves – take care of themselves at $r = 0$ and they give no contribution in relevant boundary terms that might appear above.

The compactness condition $R(r, \eta) \ll \lambda(\eta)$ means that the areal size of the radiation pulse is small comparing to the critical radius $\lambda(\eta)$. This in turn implies, using the Fourier transforms [10], that in this (no-backscatter) case the spectrum is dominated by high frequencies. The compelling explanation of the observed correlation (dominant high frequency)–(no backscatter) is that short length-waves do not undergo backscatter. If the above condition is not satisfied, then we do not have analytic arguments that would necessitate strong backscatter, but a preliminary numerical analysis suggests that this indeed is the case.

In conclusion, short gravitational waves – with the adjective "short" meaning a wavelength much smaller than the Hubble radius for the $k = 0, -1$ universes – propagate without backscatter. Longer gravitational waves, in particular those with the wavelengths being of the order of the radius of the horizon (for $k = 0, -1$ Friedman models, and also for the $k = 1$ model, under additional conditions) are expected to backscatter during the matter dominated era. This effect can manifest most strongly in the superhorizon scale.

-
- [1] S. Weinberg, *Gravitation and Cosmology*, Wiley and Sons 1972.
 - [2] J. Hadamard *Lectures on Cauchy's problem in linear partial differential equations*, Yale University Press, Yale, New Haven 1923.
 - [3] C. V. Vishveshwara, *Nature* **227**, 936(1970).
 - [4] R. Price, *Phys. Rev.* **D5**, 2419(1972).
 - [5] J. Karkowski, Z. Świerczyński and E. Malec, *Classical and Quantum Gravity* **21**, 1303(2004).
 - [6] T. Regge and J. A. Wheeler, *Phys. Rev.* **108**, 1063(1957).
 - [7] F. J. Zerilli, *Phys. Rev.* **D2**, 2141(1970).
 - [8] E. Malec, G. Wylęzek and J. Karkowski, *General Relativity and Gravitation* **36** 2151(2004).
 - [9] M. S. Berger, *Nonlinearity and functional analysis*, Academic Press, New York, San Francisco, New York 1977.
 - [10] R. Bracewell, *The Fourier Transform and its applications*, McGraw-Hill Book Company 1965.